OZPACS: Recent impacts on Australian ecosystems Reference List

Users please note: Although we endeavoured to make this list as complete and up to date as possible, some references are incomplete in cases where the full information was not available using a range of search methods. Hopefully sufficient information is provided should you wish to undertake a more thorough search for a particular reference. K. Fitzsimmons, 31/8/2007

Adamson, K., Tibby, J., Kershaw, A.P., in prep. Long term water quality and vegetation patterns at Junction Park Billabong, Murray River, Australia, with special emphasis on European impact. *Austral Ecology*.

Agnew, C.F., 2002. Recent sediment dynamics and contaminant distribution folloing a bushfire in the Nattai River catchment, NSW. Honours Thesis, University of Wollongong, Wollongong.

Aitken, D., Kershaw, A.E., 1992. Holocene vegetation and environmental history of Cranbourne Botanic Garden. *Proceedings of the Royal Society of Victoria* **105**, 67-80.

Allan, T., 2006. Examining diatoms as biological indicators of water quality, within 2 coastal lakes. School of Environmental Science and Management, Southern Cross University, Lismore.

Allen, V., Head, L., Medlin, G., Witter, D., 2000. Palaeo-ecology of the Gap and Coturaundee Ranges, western New South Wales, using stick-nest rat (Leporillus spp.) (Muridae) middens. *Austral Ecology* **25**, 333-343.

Anderson, P., 1986. A palaeoenvironmental and stratigraphic history from lowland swamp environments: Mulgrave River, north-east Queensland. Honours Thesis, Monash University, Melbourne.

Anker, S.A., Colhoun, E.A., Barton, C.E., Peterson, M., Barbetti, M., 2001. Holocene Vegetation and Paleoclimatic and Paleomagnetic History from Lake Johnston, Tasmania. *Quaternary Research* **56**, 264-274.

AQUA, 1991. Booklet for 1991 Field Meeting. Australian Quaternary Association, Mallacoota.

Ashton, D.H., Hargreaves, G.R., 1993. Dynamics of subalpine vegetation at Echo Flat, Lake Mountain, Victoria. *Proceedings of the Ecological Society of Australia* **12**, 35-60.

Atahan, P., Dodson, J.R., Itzstein-Davey, F., 2004. A fine-resolution Pliocene pollen and charcoal record from Yallalie, south-western Australia. *Journal of Biogeography* **31**, 199-205.

Atchison, J., Head, L., Fullagar, R., 2005. Archaeobotany of fruit seed processing in a monsoon savanna environment: evidence from the Keep River region, Northern territory, Australia. *Journal of Archaeological Science* **32**, 167-181.

Balme, B.E., 1963. Palynological report 98: Lake Eyre No. 20 Bore, South Australia. Geological Survey of South Australia Reports of Investigations 24, Adelaide.

Barker, A.B., 2000. Towards an archaeobotanical reference bank: a pilot study for the construction of an identification tool for macroscopic plant remains from archaeological deposits. Honours Thesis, The Flinders University of South Australia, Adelaide.

Barnett, E., 1994. A Holocene paleoenvironmental history of Lake Alexandrina, South Australia. *Journal of Paleolimnology* **12**, 259-268.

Barry, M.J., Tibby, J., Tsitsilas, A., Mason, B., Kershaw, P., Heijnis, H., 2005. A long term lake salinity record and its relationship to Daphnia populations. *Archiv für Hydrobiologie* **163**, 1-23.

Baumber, A.P., 2001. Holocene infill and evolution of Lake Wollumboola, a saline coastal lake on the New South Wales South coast. Honours Thesis, University of Wollongong, Wollongong.

Beaton, J.M., 1977. Dangerous Harvest: Investigations in the late pre-historic occupation of upland south-east central Queensland. PhD Thesis, Australian National University, Canberra.

Becker, C. PhD Thesis, University of New South Wales, Sydney.

Bell, C.J.E., Finlayson, B.L., Kershaw, A.P., 1989. Pollen analysis and dynamics of a peat deposit in Carnarvon National Park, central Queensland. *Australian Journal of Ecology* **14**, 449-456.

Berry, S.L., 1991. The potential of fossil mammal middens as indicators of vegetation history in central Australia. *Australian Journal of Botany* **39**, 305-313.

Bickford, S., Gell, P., 2005. Holocene vegetation change, Aboriginal wetland use and the impact of European settlement on the Fleurieu Peninsula, South Australia. *The Holocene* **15**, 200-215.

Binder, R.M., 1978. Stratigraphy and pollen analysis of a peat deposit, Bunyip Bog, Mt Buffalo, Victoria. Department of Geography, Monash University, Melbourne.

Binder, R.M., Kershaw, A.P., 1978. A Late Quaternary pollen diagram from the southeastern highlands of Australia. *Search* **9**, 44-45.

Black, M., 2001. Fire, vegetation, humans and climate: A record of change during the past millennium from the King's Tableland Swamp, Blue Mountains, New South Wales. Honours Thesis, University of New South Wales, Sydney.

Black, M.P., 2006. A Late Quaternary palaeoenvironmental investigation of the fire, climate, human and vegetation nexus from the Sydney Basin, Australia. PhD Thesis, University of New South Wales, Sydney.

Black, M.P., Mooney, S.D., 2006. Holocene fire history from the Greater Blue Mountains World Heritage Area, New South Wales, Australia: the climate, humans and fire nexus. *Regional Environmental Change* **6**, 41-51.

Black, M.P., Mooney, S.D., 2007. The response of Aboriginal burning practices to population levels and El Nino-Southern Oscillation events during the mid- to late-Holocene: a case study from the Sydney Basin using charcoal and pollen analysis. *Australian Geographer* **38**, 37-52.

Black, M.P., Mooney, S.D., Martin, H.A., 2006. A >43,000-year vegetation and fire history from Lake Baraba, New South Wales, Australia. *Quaternary Science Reviews* **25**, 3003-3016.

Bohte, A., 1994. Local Vegetation and Environments of the Last Glacial Cycle at Lynch's Crater, Northeast Queensland. Honours Thesis, Monash University, Melbourne.

Bohte, A., Kershaw, A.P., 1999. Taphonomic influences on the interpretation of the palaeoecological record of Lynch's Crater, northeastern Australia. *Quaternary International* **57-58**, 49-59.

Boon, S., Dodson, J.R., 1992. Environmental response to land use at Lake Curlip, east Gippsland, Victoria. *Australian Geographical Studies* **30**, 206-221.

Bottomley, C., 1994. Late Quaternary palaeoenvironments of Carlisle Perched Lake in the Otway Ranges, Victoria. Honours Thesis, Monash University, Melbourne.

Bowler, J.M., Hamada, T., 1971. Late Quaternary stratigraphy and radiocarbon chronology of water level fluctuations in Lake Keilambete, Victoria. *Nature* **232**, 330-332.

Bowler, J.M., Qi, H., Kezao, C., Head, M.J., Baoyin, Y., 1986. Radiocarbon dating of playa-lake hydrologic changes: examples from northwestern China and central Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **54**, 241-260.

Boyd, W.E., 1990. Quaternary pollen analysis in the arid zone of Australia: Dalhousie Springs, central Australia. *Review of Palaeobotany and Palynology* **64**, 331-341.

Boyd, W.E., 1994. Quaternary pollen analysis in the Arid Zone of Australia: Further Results from Dalhousie Springs, central Australia. *Australian Geographical Studies* **32**, 274-280.

Boyd, W.E., 1994. Research Note: Quaternary Pollen Analysis in the Arid Zone of Australia. *Australian Geographical Studies* **32**, 274-280.

Boyd, W.E., Lentfer, C.J., Parr, J.F., 2005. Interactions between human activity, volcanic eruptions and vegetation during the Holocene at Garua and Numundo, West New Britain, PNG. *Quaternary Research* **64**, 384-398.

Boyd, W.E., Lentfer, C.J., Parr, J.F., Jago, L.C.F., 2004. Holocene vegetation dynamics in West New Britain, PNG, International Palynological Conference, Granada.

Boyd, W.E., Stubbs, B.J., Averill, C., 1999. The 'grasses' of the Big Scrub District of north-eastern New South Wales: a sedimentary record of late Holocene grasslands in subtropical forest landscape. *Australian Geographer* **30**, 331-336.

Brazier, J., in prep. Fate of Heavy metal contaminants from Rum Jungle Uranium Mine into the Finniss River, Northern Territory, Australia. PhD Thesis, Charles Darwin University, Darwin.

Buchanan, G., 2001. Chronology of sediment and land-use/vegetation history in the Kakadu area, NT. Honours Thesis, University of Technology, Sydney, Sydney.

Cameron, N.G., Tyler, P.A., Rose, N.L., Hutchinson, S., Appleby, P.G., 1993. The recent palaeolimnology of Lake Nicholls, Mount Field National Park, Tasmania. *Hydrobiologia* **269-270**, 361-370.

Cassidy, H., 2001. The water quality history of Lake Hiawatha, northern NSW, using diatoms as bioindicators. School of Environmental Science and Management, Southern Cross University, Lismore.

Cavanagh, J.E., Burns, K.A., Brunskill, G.J., Coventry, R.J., 1999. Organochlorine pesticide residues in soils and sediments of the Herbert and Burdekin River regions, North Queensland - implications for contamination of the Great Barrier Reef. *Marine Pollution Bulletin* **39**, 367-375.

Cavanagh, J.E., Burns, K.A., Brunskill, G.J., Coventry, R.J., 2003. Organochlorine insecticide usage in the sugar industry of the Herbert and Burdekin River regions: chemical and biological assessments. CRC for Sustainable Sugar Production, Townsville.

Chalson, J.M., 1983. Honours Thesis, University of New South Wales, Sydney.

Chalson, J.M., 1991. The late Quaternary climatic and vegetation history of the Blue Mountains, NSW. PhD Thesis, University of New South Wales, Sydney.

Chivas, A.R., García, A., van der Kaars, S., Couapel, M.J.J., Holt, S., Reeves, J.M., Wheeler, D.J., Switzer, A.D., Murray-Wallace, C.V., Banerjee, D., Price, D.M., Wang, S.X., Pearson, G., Edgar, N.T., Beaufort, L., De Deckker, P., Lawson, E., Cecil, C.B., 2001. Sea-level and environmental changes since the Last Interglacial in the Gulf of Carpentaria. *Quaternary International* 83-85, 19-46.

Churchill, D.M., 1959. Late Quaternary eustatic changes in the Swan River District. *Journal of the Royal Society of Western Australia* **42**, 53-55.

Churchill, D.M., 1960. Late Quaternary changes in the vegetation on Rottnest Island. *Western Australian Naturalist* **7**, 160-166.

Churchill, D.M., 1968. The distribution and prehistory of Eucalypt diversicolor F. Muell., E. marginata Donn ex Sm., and E. calophylla R. Br. in relation to rainfall. *Australian Journal of Botany* **16**.

Churchill, D.M., Dodson, J.R., 1980. The Occurrence of Phyllocladus aspleniifolius (Labill.) Hook.F. in Victoria prior to 1100B.P. *Muelleria* **4**, 277-284.

Churchill, D.M., Galloway, R.W., Singh, G., 1978. Closed lakes and the palaeoclimatic record. In: Pittock, A.B., Frakes, L.A., Jenssen, D., Peterson, J.A., Zillman, J.W. (Eds.), Climate Change and Variability: a Southern Hemisphere Perspective. Cambridge University Press, Cambridge, pp. 97-108.

Clark, 1997. Honours Thesis.

Clark, R.L., 1976. Vegetation history and the influence of the sea at Lashmar's Lagoon, Kangaroo Island, South Australia, 3,000 B.P. to the Present Day. Honours Thesis, Monash University, Melbourne.

Clark, R.L., 1983. Pollen and charcoal evidence for the effects of Aboriginal burning on the vegetation of Australia. *Archaeology in Oceania* **18**, 32-37.

Clark, R.L., 1986. The fire history of Rotten Swamp, A.C.T. Unpublished Report to ACT Parks and Conservation Service, CSIRO, Canberra.

Clark, R.L., 1990. Ecological history for environmental management. *Proceedings of the Ecological Society of Australia* **16**, 1-16.

Clark, R.L., Guppy, J.C., 1988. A transition from mangrove forest to freshwater wetland in the monsoon tropics of Australia. *Journal of Biogeography* **15**, 665-684.

Clark, R.L., Hope, G.S., Hope, J.H., in press. Palaeoenvironments of the Rocky River bone bed, Kangaroo Island, South Australia. *Transactions of the Royal Society of South Australia*.

Clarke, A., 1985. A preliminary archaeobotanical analysis of the Anbangbang 1 site. In: Jones, R. (Ed.), Archaeological Research in Kakadu National Park. Australian National Parks and Wildlife Service, pp. 77-96.

Clarke, A., 1988. Archaeological and ethnobotanical interpretations of plant remains from Kakadu National Park, Northern Territory. In: Meehan, B., Jones, R. (Eds.), Archaeology with ethnography: an Australian perspective. Department of Prehistory, Australian National University, Canberra.

Clarke, A., 1989. Macroscopic plant remains. In: Beck, W., Clarke, A., Head, L. (Eds.), Plants in Australian Archaeology. TEMPUS.

Coddington, J., 1983. Landscape evolution and its effect on Aboriginal occupation at Tilba Tilba, south coast New South Wales. Honours Thesis, Australian National University, Canberra.

Colhoun, E., et al., 1991. Australian Geographical Studies 29, 337-354.

Colhoun, E., Goede, A., 1974. A Reconnaissance Survey of the Glaciation of Macquarie Island. *Proceedings of the Royal Society of Tasmania* **108**, 1-19.

Colhoun, E.A., 1977. The Remarkable Cave, Southeastern Tasmania: its geomorphogical development and environmental history. *Papers and Proceedings of the Royal Society of Tasmania* **111**, 29-39.

Colhoun, E.A., 1985. Pre-last glaciation maximum vegetation history at Henty Bridge, Western Australia. *New Phytologist* **100**, 681-690.

Colhoun, E.A., 1992. Late Glacial and Holocene vegetation history at Poets Hill Lake, Western Tasmania. *Australian Geographer* **23**, 11-23.

Colhoun, E.A., 1996. Application of Iversen's glacial-interglacial cycle to interpretation of the late last glacial and holocene vegetation history of western Tasmania. *Quaternary Science Reviews* **15**, 557-580.

Colhoun, E.A., Pola, J.S., Barton, C.E., Heijnis, H., 1999. Late-Pleistocene vegetation and climate history of Lake Selina, western Tasmania. *Quaternary International* **57-58**, 5-23.

Colhoun, E.A., Sigleo, W.R., 1981. A short pollen diagram from Crown Lagoon in the Midlands of Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* **115**, 181-188.

Colhoun, E.A., van de Geer, G., 1987. Holocene to Middle Last Glaciation vegetation history at Tullabardine Dam, western Tasmania. *Proceedings of the Royal Society of London Series B* **229**, 177-207.

Colhoun, E.A., van de Geer, G., 1987. Vegetation history and climate before the Maximum of the Last Glaciation at Crotty, Western Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* **121**, 69-74.

Colhoun, E.A., van de Geer, G., 1988. Darwin Crater, the King and Linda Valleys. In: Colhoun, E.A. (Ed.), Cainozoic vegetation in Tasmania. University of Newcastle, Newcastle, pp. 30-71.

Colhoun, E.A., van de Geer, G., Hannan, D., 1991. Late Glacial and Holocene vegetation history at Dublin Bog, north-central Tasmania. *Australian Geographical Studies* **29**, 337-354.

Colhoun, E.A., van de Geer, G., Mook, W.G., 1982. Stratigraphy, pollen analysis and paleoclimatic interpretation of Pulbeena Swamp, northwestern Tasmania. *Quaternary Research* **18**, 108-126.

Colliton, J.P., 2001. The impact of European settlement and coal mining on the Nattai Catchment, NSW. Honours Thesis, University of Technology, Sydney, Sydney.

Cook, D.E., 2006. A 2000 year record of environmental change from Tocal Homestead Lagoon, eastern Australia. PhD Thesis, The University of Sydney, Sydney.

Costin, A.B., 1972. Carbon-14 dates from the Snowy Mountains area, southeastern Australia, and their interpretation. *Quaternary Research* **2**, 579-590.

Croome, R., Kristiansen, J., Tyler, P., 1998. A Description of Mallomonas Marsupials nov. sp. (Synurophyceae), a new chrysophyte from Australia, with comments on the endemicity of Australian freshwater algae. *Nordic Journal of Botany* **18**, 633-639.

Crowley, G.M., 1981. The Late Quaternary environmental history of the Lake Bolac region of western Victoria, and its implications for Aboriginal occupation. Honours Thesis, Monash University, Melbourne.

Crowley, G.M., 1994. Quaternary soil salinity events and Australian vegetation history. *Quaternary Science Reviews* **13**, 15-22.

Crowley, G.M., Anderson, P., Kershaw, A.P., Grindrod, J.F., 1990. Palynology of a Holocene marine transgressive sequence, lower Mulgrave River valley, north-east Queensland. Australian *Journal of Ecology* **15**.

Crowley, G.M., Gagan, M.K., 1995. Holocene evolution of coastal wetlands in wet - tropical Australia. *The Holocene* **5**, 385-399.

Crowley, G.M., Grindrod, J., Kershaw, A.P., 1994. Modern pollen deposition in the tropical lowlands of north east Queensland, Australia. *Review of Palaeobotany and Palynology* **83**, 299-327.

Crowley, G.M., Kershaw, A.P., 1994. Late Quaternary environmental change and human impact around Lake Bolac, western Victoria, Australia. *Journal of Quaternary Science* **9**, 367-377.

Cupper, M.L., Drinnan, A.N., Thomas, I., 2000. Holocene palaeoenvironments of salt lakes in the Darling Anabranch region, south-western New South Wales, Australia. *Journal of Biogeography* **27**, 1079-1094.

D'Costa, D.M., Kershaw, A.P., 1997. An Expanded Recent Pollen Database from South-eastern Australia and its Potential for Refinement of Palaeoclimatic Estimates. *Australian Journal of Botany* **45**, 583-605.

D'Costa, D.M.M., 1989. Late Quaternary vegetation and environments from Lake Terang, Western Australia. MSc Thesis, Monash University, Melbourne.

D'Costa, D., 1997. The reconstruction of Quaternary vegetation and climate on King Island, Bass Strait, Australia. PhD Thesis, Monash University, Melbourne.

D'Costa, D.M., Edney, P., Kershaw, P., De Deckker, P., 1989. Late Quaternary palaeoecology of Tower Hill, Victoria, Australia. *Journal of Biogeography* **16**, 461-482.

D'Costa, D.M., Grindrod, J., Ogden, R., 1993. Preliminary environmental reconstructions from late Quaternary pollen and mollusc assemblages at Egg Lagoon, King Island, Bass Strait. *Australian Journal of Ecology* **18**, 351-366.

De Deckker, P., 1982. Holocene Ostracods, other invertebrates and fish remains from cores of four maar lakes in southeastern Australia. *Proceedings of the Royal Society of Victoria* **94**, 183-220.

Dimitriadis, S., Cranston, P.S., 2001. An Australian Holocene climate reconstruction using Chironomidae from a tropical volcanic maar lake. *Palaeogeography, Palaeoclimatology, Palaeoecology* **176**, 109-131.

Dodson, J.R., 1974. Vegetation and climatic history near Lake Keilambete, western Victoria. *Australian Journal of Botany* **22**, 709-717.

Dodson, J.R., 1974. Vegetation history and water fluctuations at Lake Leake, South-eastern South Australia Part 1. 10,000 B.P. to present. *Australian Journal of Botany* **22**, 719-741.

Dodson, J.R., 1975. Late Quaternary vegetation history of southeastern South Australia. PhD Thesis, Australian National University, Canberra.

Dodson, J.R., 1975. The pre-Settlement vegetation of the Mt. Gambier area, South Australia. *Transactions of the Royal Society of South Australia* **99**, 89-92.

Dodson, J.R., 1975. Vegetation history and water fluctuations at Lake Leake, south-eastern South Australia. Part 2. 50,000 BP to 10,000 BP. *Australian Journal of Botany* **23**, 815-831.

Dodson, J.R., 1977. Pollen deposition in a small closed drainage basin lake. *Review of Palaeobotany and Palynology* **24**, 179-193.

Dodson, J.R., 1982. Modern pollen rain and recent vegetation history on Lord Howe Island: Evidence of human impact. *Review of Paleobotany and Palynology* **38**, 1-21.

Dodson, J.R., 1982. Modern pollen rain in southeastern New South Wales, Australia. *Review of Palaeobotany and Palynology* **38**, 249-268.

Dodson, J.R., 1983. Pollen recovery from organic lake clays: A comparison of two techniques. *Pollen et Spores* **XXV**, 131-138.

Dodson, J.R., 1986. Holocene vegetation and environments near Goulburn, New South Wales. *Australian Journal of Botany* **34**, 231-249.

Dodson, J.R., 1987. Mire Development and Environmental Change, Barrington Tops, New South Wales, Australia. *Quaternary Research* **27**, 73-81.

Dodson, J.R., 1988. The perspective of pollen records to study response, competition and resilience in vegetation on Barrington Tops, Australia. *Progress in Physical Geography* **12**, 183-208.

Dodson, J.R., 2000. Radiocarbon dates from a Holocene deposit in Southwestern Australia. *Radiocarbon* **42**, 229-234.

Dodson, J.R., 2001. A vegetation and fire history in a subalpine woodland and rain-forest region, Solomons Jewel Lake, Tasmania. *The Holocene* **11**, 111-116.

Dodson, J.R., Chant, J., Daly, J., 1995. Human impact recorded in urban wetlands sediments in Sydney, Australia. *Man and Culture in Oceania* **11**, 113-124.

Dodson, J.R., De Salis, I., Myers, C.A., Sharp, A.J., 1994. A thousand years of environmental change and human impact in the alpine zone at Mount Kosciuszko, New South Wales. *Australian Geographer* **25**, 77-87.

- Dodson, J.R., Frank, K., Fromme, M., Hickson, D., McRae, Mooney, S., Smith, J.D., 1994. Environmental Systems and Human Impact at Cobrico Crater, South Western Victoria. *Australian Geographical Studies* **32**, 27-40.
- Dodson, J.R., Greenwood, P.G., Jones, R.L., 1986. Holocene forest and wetland dynamics at Barrington Tops, NSW. *Journal of Biogeography* **13**, 561-585.
- Dodson, J.R., Lu, J.J., 2000. A late Holocene vegetation and environment record from Byenup Lagoon, south-western Australia. *Australian Geographer* **31**, 41-54.
- Dodson, J.R., McRae, V.M., Molloy, K., Roberts, F., Smith, J.D., 1993. Late Holocene human impact on two coastal environments in New South Wales, Australia: a comparison of Aboriginal and European impacts. *Vegetation History and Archaeobotany* 2, 89-100.
- Dodson, J.R., Mitchell, F.J.G., Bogeholz, H., Julian, N., 1998. Dynamics of temperate rainforest from fine resolution pollen analysis, upper Ringarooma river, northeastern Tasmania. *Australian Journal of Ecology* **23**, 550-561.
- Dodson, J.R., Mooney, S.D., 2002. An assessment of historic human impact on south-eastern Australian environmental systems, using late Holocene rates of environmental change. *Australian Journal of Botany* **50**, 455-464.
- Dodson, J.R., Roberts, F.K., De Salis, T., 1994. Palaeoenvironments and human impact at Burraga Swamp in montane rainforest, Barrington Tops National Park, New South Wales, Australia. *Australian Geographer* **25**, 161-169.
- Dodson, J.R., Robinson, M., Tardy, C., 2005. Two fine-resolution Pliocene charcoal records and their bearing on pre-human fire frequency in south-western Australia. *Austral Ecology* **30**, 592-599.
- Dodson, J.R., Thom, B.G., 1992. Holocene vegetation history from the Hawkesbury Valley, New South Wales. *Proceedings of the Linnean Society of New South Wales* **113**, 121-134.
- Donders, T.H., Haberle, S.G., Hope, G.S., Wagner, F., Visscher, H., 2007. Pollen evidence for the transition of the Eastern Australian climate system from the post-glacial to the present-day ENSO mode. *Quaternary Science Reviews* **26**, 1621-1637.
- Donders, T.H., Wagner, F., Visscher, H., 2006. Late Pleistocene and Holocene subtropical vegetation dynamics recorded in perched lake deposits on Fraser Island, Queensland, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **241**, 417-439.
- Dyson, W.D., 1995. A pollen and vegetation history from Lake Dove, Western Tasmania. Honours Thesis, University of Newcastle, Newcastle.
- Edgar, G.J., Samson, C.R., 2004. Catastrophic decline in mollusc diversity in Eastern Tasmania and its concurrence with shellfish fisheries. *Conservation Biology* **18**, 1579-1588.
- Edney, P., Kershaw, A.P., Peterson, J.A., Coutts, P.J.F., 1985. Evidence of a Pleistocene age for Tower Hill, western Victoria. *Search* **16**, 302-303.

- Edney, P.A., Kershaw, A.P., De Deckker, P., 1990. A late Pleistocene and Holocene vegetation and environmental record from Lake Wangoom, Western Plains of Victoria, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **80**, 325-343.
- Edwards, S., McKirdy, D., Bone, Y., Gell, P., Gostin, V.A., 2006. Diatoms and ostracods as mid-Holocene paleoenvironmental indicators, North Stromatolite Lake, Coorong National Park, South Australia. *Australian Journal of Earth Sciences* **53**, 651-663.
- Ellis, R.C., Thomas, I., 1988. Pre-settlement and post-settlement vegetational change and probable Aboriginal influences in a highland forested area in Tasmania. In: Frawley, K.J., Semple, N.M. (Eds.), Australia's Ever-changing Forests: Proceedings of the First National Conference on Australian Forest History. Department of Geography and Oceanography, Australian Defence Force Academy, Canberra.
- Esteban, G., Finlay, B., Olmo, J., Tyler, P., 2000. Ciliated protozoa from a volcanic crater-lake in Victoria, Australia. *Journal of Natural History* **34**, 159-189.
- Farago, L., 2001. Using diatoms as bioindicators of past and contemporary water quality conditions in Tuckean Swamp, northeastern NSW. School of Environmental Science and Management, Southern Cross University, Lismore.
- Farago, L., 2002. Using diatoms and pollen as bioindicators of past and contemporary environmental conditions in Tuckean Swamp, Northeastern NSW. Honours Thesis, Southern Cross University, Lismore.
- Field, J.H., Dodson, J.R., Prosser, I.P., 2002. A late Pleistocene vegetation history from the Australian semi-arid zone. *Quaternary Science Reviews* **21**, 1023-1037.
- Fitzgerald, F., 1998. The impact of European settlement on two lagoons in New South Wales, Australia. Masters Thesis, University of Wollongong, Wollongong.
- Fitzsimmons, K.E., Rhodes, E.J., Magee, J.W., Barrows, T.T., in press. The timing of linear dune activity in the Strzelecki and Tirari Deserts, Australia. *Quaternary Science Reviews*.
- Fitzsimons, S.J., Colhoun, E.A., van de Geer, G., Roberts, S., 1990. Definition and character of the Regency Interglacial and Early-Middle Pleistocene stratigraphy in the King Valley, western Tasmania, Australia. *Boreas* **19**, 1-15.
- Flannery, T.F., Gott, B., 1984. The Spring Creek locality, southwestern Victoria, a late surviving megafaunal assemblage. *Australian Zoologist* **21**, 385-422.
- Flett, I., 2003. The history of algal blooms in the Myall Lakes. Honours Thesis, University of Technology, Sydney, Sydney.
- Fluin, J., Gell, P., Haynes, D., Tibby, J., submitted. Paleolimnological evidence for the independent evolution of neighbouring terminal lakes, the Murray Darling Basin, Australia. *Hydrobiologia*.

- Gagan, M.K., Johnson, D.D., Crowley, G.M., 1994. Sea level control of stacked late Quaternary coastal sequences, central Great Barrier Reef. *Sedimentology* **41**, 329-351.
- Gale, S.J., 2003. Making the European landscape: early contact environmental impact in Australia, Geography's New Frontiers: Geographical Society of New South Wales Conference. Geographical Society of New South Wales, University of New South Wales, Sydney.
- Gale, S.J., Cook, D.E., 2006. The 210Pb chronology of deposition in Tocal Homestead Lagoon, eastern Australia. *Quaternary Newsletter* **110**, 40-43.
- Gale, S.J., Haworth, R.J., 2002. Beyond the Limits of Location: human environmental disturbance prior to official European contact in early colonial Australia. *Archaeology in Oceania* **37**, 123-136.
- Gale, S.J., Haworth, R.J., 2005. Catchment-wide soil loss from pre-agricultural times to the present: transport- and supply-limitation of erosion. *Geomorphology* **68**, 314-333.
- Gale, S.J., Haworth, R.J., Cook, D.E., Williams, N.J., 2004. Human impact on the natural environment in early colonial Australia. *Archaeology in Oceania* **39**, 148-156.
- Gale, S.J., Haworth, R.J., Pisanu, P.C., 1995. The 210Pb chronology of late Holocene deposition in an eastern Australian lake basin. *Quaternary Science Reviews* **14**, 395-408.
- Gale, S.J., Pisanu, P.C., 2001. The late-Holocene decline of Casuarinaceae in southeast Australia. *The Holocene* **11**, 485-490.
- García, A., Chivas, A.R., 2004. The euryhaline genus Lamprothamnium (Charales, Charophyta) from Australia: statistical analyses and application to paleoenvironmental reconstruction. *Journal of Paleolimnology* **31**, 321-341.
- García, A., Chivas, A.R., 2006. Diversity and ecology of extant and Quaternary Australian charophytes (Charales). *Cryptogamie Algologie* **27**, 323-340.
- García, A., Jones, B.G., Chenhall, B.E., Murray-Wallace, C., 2002. Charophytes as environmental indicators: a Holocene example from Tom Thumbs Lagoon, NSW. *Alcheringa* **26**, 507-518.
- Gell, P., Baldwin, D., Little, F., Tibby, J., Hancock, G., submitted. The impact of regulation and salinisation on floodplain lakes: the lower River Murray, Australia. *Hydrobiologia*.
- Gell, P., Bulpin, S., Wallbrink, P., Bickford, S., Hancock, G., 2005. Tareena Billabong A palaeolimnological history of an everchanging wetland, Chowilla Floodplain, lower Murray-Darling Basin. *Marine and Freshwater Research* **56**, 441-456.
- Gell, P., Fluin, J., Tibby, J., Haynes, D., Khanum, S., Walsh, B., Hancock, G., Harrison, J., Zawadzki, A., Little, F., 2006. Changing fluxes of sediments and salts as recorded in lower River Murray wetlands, Australia. In: Rowan, J.S., Duck, R.W., Werritty, A. (Eds.), Sediment Dynamics and the Hydromorphology of Fluvial Systems (Proceedings of a symposium held in Dundee, UK, July 2006). International Association of Hydrological Sciences, United Kingdom, pp. 416-424.

- Gell, P., Little, F., submitted. Long Term Water Quality Changes in Murrumbidgee Floodplain Wetlands Revealed by Fossil Diatom Assemblages. *Australian Journal of Ecology*.
- Gell, P., Tibby, J., Fluin, J., Leahy, P., Reid, M., Adamson, K., Bulpin, S., MacGregor, A., Wallbrink, P., Hancock, G., Walsh, B., 2005. Accessing limnological change and variability using fossil diatom assemblages, south-east Australia. *River Research and Applications* **21**, 257 269.
- Gell, P., Wallbrink, P., Tassicker, G., Illman, M., 1999. Secrets in the sediments: a history of sediment and pollution loads in the Lower Torrens River, S.A. In: Rutherfurd, I., Bartley, R. (Eds.), Second Australian Stream Management Conference: The Challenge of Rehabilitating Australia's Streams. CRC Catchment Hydrology, Melbourne.
- Gell, P.A., 1998. Quantitative reconstructions of the Holocene palaeosalinity of paired crater lakes based on a diatom transfer function. *Palaeoclimates Data and Modelling* **4**, 1-14.
- Gell, P.A., Barker, P.A., De Deckker, P., Last, W.M., Jellad, L., 1994. The Holocene history of West Basin Lake, Victoria, Australia, chemical changes based on fossil biota and sediment mineralogy. *Journal of Palaeolimnology* **12**, 235-258.
- Gell, P.A., Stuart, I.M., 1989. Human settlement history and environmental impact: the Delegate River catchment, East Gippsland, Victoria. Monash Publications in Geography 36, Melbourne.
- Gell, P.A., Stuart, I.M., Smith, J.D., 1993. The response of vegetation to changing fire regimes and human activity in the Delegate River catchment, East Gippsland, Victoria. *The Holocene* **3**, 150-160.
- Genever, M., Grindrod, J., Barker, B., 2003. Holocene palynology of Whitehaven Swamp, Whitsunday Island, Queensland, and implications for the regional archaeological record. *Palaeogeography, Palaeoclimatology, Palaeoecology* **201**, 141-156.
- Gerritse, R.G., Hernandez, F., Murray, A.S., Wallbrink, P.J., Brunskill, G., 1995. Distribution of Organochlorines, Polycyclic Aromatic Hydrocarbons, Phosphorus and 137Cs in Sediment Profiles from Ellen Brook in Western Australia. *Marine and Freshwater Research* 46, 843-851.
- Gerritse, R.G., Wallbrink, P.J., Murray, A.S., 1998. Accumulation of Phosphorus and Heavy Metals in the Swan-Canning Estuary, Western Australia. *Estuarine, Coastal and Shelf Science* **47**, 165–179.
- Gillespie, R., Horton, D.R., Ladd, P.G., Macumber, P.G., Rich, T.H., Wright, R.V.S., 1978. Lancefield Swamp and the extinction of the Australian megafauna. *Science* **200**, 1044-1048.
- Goodfield, M.L., 1983. A Holocene pollen analytical study of Lake Eacham, Atherton Tableland, north-east Queensland. Honours Thesis, Monash University, Melbourne.
- Green, D., Singh, G., Pollach, H., Moss, D., Banks, J., Geissler, E.A., 1988. A fine resolution palaeoecology and palaeoclimatology from south-eastern Australia. *Journal of Ecology* **76**, 790-806.

Grindrod, J., 1979. Frozen sediments from a tropical lake: a palaeoecological study of Lake Eacham, north-east Queensland. Honours Thesis, Monash University, Melbourne.

Grindrod, J., 1985. The palynology of mangroves on a prograded shore, Princess Charlotte Bay, North Queensland, Australia. *Journal of Biogeography* **12**, 323-348.

Grindrod, J., 1988. Holocene mangrove history of the South Alligator River estuary, Northern Territory, Australia. PhD Thesis, Australian National University, Canberra.

Grindrod, J., 1988. The palynology of Holocene mangrove and saltmarsh sediments, particularly in Northern Australia. *Review of Palaeobotany and Palynology* **55**, 229-245.

Grindrod, J., Rhodes, E.G., 1984. Holocene sea-level history of a tropical estuary: Missionary Bay, North Queensland. In: Thom, B.G. (Ed.), Coastal Geomorphology in Australia. Academic Press, New York, pp. 151-178.

Groves, 2001. Honours Thesis, University of Western Australia, Perth.

Haberle, S.G., 2005. A 23,000-yr pollen record from Lake Euramoo, Wet Tropics of NE Queensland, Australia. *Quaternary Research* **64**, 343-356.

Haberle, S.G., Tibby, J.C., Dimitriadis, S., Heijnis, H., 2006. The impact of European occupation on terrestrial and aquatic ecosystem dynamics in an Australian tropical rain forest. *Journal of Ecology* **94**, 987-1002.

Hancock, G., Pietsch, T., 2006. Sedimentation in the Gippsland Lakes as determined from sediment cores. CSIRO Land and Water Science Report 40/06, Canberra.

Hancock, G.J., 2001. Sediment accumulation in central Moreton Bay as determined from sediment core profiles. Report on Sediment Source Project Phase 3, Part A. CSIRO Land and Water, Canberra.

Hancock, G.J., Hunter, J.R., 1999. Use of excess 210Pb and 228Th to estimate rates of sediment accumulation and bioturbation in Port Phillip Bay, Australia. *Marine and Freshwater Research* **50**, 533-545.

Hancock, G.J., Olley, J.M., Wallbrink, P.J., 2001. Sediment transport and accumulation in Western Port: Report on Phase 1 of a study determining the sources of sediment to Western Port. CSIRO Land and Water Technical Report 47/01, Canberra.

Hargreaves, G.R., 1977. Vegetation dynamics of Echo Flat, Lake Mountain, Victoria. MSc Thesis, University of Melbourne, Melbourne.

Harle, K., 1989. Palaeoenvironments of the Mt Anne region in south western Tasmania. Honours Thesis, Monash University, Melbourne.

- Harle, K.J., 1997. Late Quaternary vegetation and climate change in southeastern Australia: palynological evidence from marine core E55-6. *Palaeogeography, Palaeoclimatology, Palaeoecology* **131**, 465-483.
- Harle, K.J., Britton, K., Heijnis, H., Zawadzki, A., Jenkinson, A.V., 2002. Mud, mines and rainforests: a short history of human impact in western Tasmania, using pollen, trace metals and lead-210. *Australian Journal of Botany* **50**, 481-497.
- Harle, K.J., Hodgson, D.A., Tyler, P.A., 1999. Palynological evidence for Holocene palaeoenvironments from the lower Gordon River valley, in the World Heritage Area of southwest Tasmania. *The Holocene* **9**, 149-162.
- Harle, K.J., Kershaw, A.P., Heijnis, H., 1999. The contributions of marine palynology and uranium/thorium to the dating of the lake Wangoom pollen record., Western Plains of Victoria, Australia. *Quaternary International* **57-58**, 25-34.
- Harrison, J., 2000. The spatial and temporal pollution distribution in the Tonalli River, NSW. Honours Thesis, University of Technology Sydney, Sydney.
- Harrison, J., H., Caprarelli, G., 2003. Historical pollution variability from abandoned mine sites, Greater Blue Mountains World Heritage Area, NSW, Australia. *Environmental Geology* **43**, 680-687.
- Hart, B., 2005. Using diatoms to assess the past and present water quality of Salty Lagoon, Evans Head. School of Environmental Science and Management, Southern Cross University, Lismore.
- Hartney, L., 2005. A Late Quaternary Vegetation History from the Archaeologically Significant Mt Eccles Lava Flow, South-west Victoria. Honours Thesis, Monash University, Melbourne.
- Hastings, 2000. Honours Thesis, University of New South Wales, Sydney.
- Haworth, R.J., 1994. European impact on lake sedimentation in upland eastern Australia: case studies from the New England Tablelands of New South Wales. PhD Thesis, The University of New England, Armidale.
- Haworth, R.J., Gale, S.J., Short, S.A., Heijnis, H., 1999. Land use and lake sedimentation on the New England Tablelands of New South Wales, Australia. *Australian Geographer* **30**, 51-73.
- Haynes, D., Gell, P., Tibby, J., Hancock, G., Goonan, P., submitted. Against the tide: the freshening of naturally saline coastal lakes, south east South Australia. *Hydrobiologia*.
- Head, L., 1983. Environment as artefact: a geographic perspective on the Holocene occupation of Southwestern Victoria. *Archaeology in Oceania* **18**, 73-80.
- Head, L., 1985. Further discussion of Phyllocladus and its pollen in Victoria during the Holocene. *Australian Journal of Ecology* **10**, 73-76.

- Head, L., 1985. Pollen analysis of sediments from the Bridgewater Caves archaeological site, southwestern Victoria. *Australian Archaeology* **20**, 1-15.
- Head, L., 1987. The Holocene prehistory of a coastal wetland system: Discovery Bay, southeastern Australia. *Human Ecology* **15**, 435-462.
- Head, L., 1988. Holocene vegetation, fire and environmental history of the Discovery Bay region, south-western Victoria. *Australian Journal of Ecology* **13**, 21-49.
- Head, L., 1989. Prehistoric Aboriginal impacts on Australian vegetation: an assessment of the evidence. *Australian Geographer* **20**, 37-46.
- Head, L., 1989. Using palaeoecology to date Aboriginal fishtraps at Lake Condah, Victoria. *Archaeology in Oceania* **24**, 110-115.
- Head, L., Stewart, I.-M.F., 1980. Change in the Aire: Palaeoecology and Prehistory in the Aire Basin, southwestern Victoria. Monash University, Melbourne.
- Hill, R.S., Macphail, M.K., 1985. A fossil flora from rafted Plio-Pleistocene mudstones at Regatta Point, Tasmaina. *Australian Journal of Botany* **33**, 497-517.
- Hodgson, D., Tyler, P., 1996. The impact of a hydroelectric dam on the stability of meromictic lakes in southwest Tasmania, Australia. *Archiv für Hydrobiologie* **137**, 301-323.
- Hodgson, D., Tyler, P.A., Vyverman, W., 1997. The palaeolimnology of Lake Fidler, a meromictic lake in south-west Tasmania and the significance of recent human impact. *Journal of Paleolimnology* **18**, 313-333.
- Hodgson, D., Vyverman, W., Chepstow-Lusty, A., Tyler, P.A., 2000. From rainforest to wasteland in 100 years The limnological legacy of the Queenstown mines, SW Tasmania. *Archiv für Hydrobiologie* **149**, 25-36.
- Hodgson, D., Vyverman, W., Tyler, P.A., 1997. Diatoms of meromictic lakes adjacent to the Gordon River, and of the Gordon River estuary in south-west Tasmania. *Bibliotheca Diatomologica* **35**, 1-173.
- Hooley, A.D., Southern, W., Kershaw, A.P., 1980. Holocene vegetation and environments of Sperm Whale Head, Victoria, Australia. *Journal of Biogeography* **7**, 349-362.
- Hope, G., 1999. Vegetation and fire response to late Holocene human occupation in island and mainland north west Tasmania. *Quaternary International* **59**, 47-60.
- Hope, G., Kershaw, P., van der Kaars, S., Ziangjun, S., Liew, P.-M., Heusser, L., .E., Takahara, H., McGlone, M., Miyoshi, N., Moss, P.T., 2004. History of vegetation and habitat change in the Australia-Asia region. *Quaternary International* **118-119**, 103-126.
- Hope, G.S., 1974. The Vegetation History from 6,000 B.P. to Present of Wilsons Promontory, Victoria, Australia. *New Phytologist* **73**, 1035-1053.

- Hope, G.S., 1978. The Late Pleistocene and Holocene Vegetational History of Hunter Island, North-Western Tasmania. *Australian Journal of Botany* **26**, 493-514.
- Hope, G.S., 1996. A report on the environment of Mt Scabby and palaeoecological analyses of swamp and stone arrangement sediments. Unpublished report to ACT National Parks and Wildlife Service, Canberra.
- Hope, G.S., Coddington, J., O'Dea, D., in press. Estuarine development and human occupation at Bobundara Swamp, Tilba Tilba, New South Wales, Australia. In: Lillie, M. (Ed.), Wetlands Archaeology and Environments: Regional Issues, Global Perspective. Routledge, London.
- Hope, G.S., Hughes, P.J., Russell-Smith, J., 1985. Geomorphological fieldwork and the evolution of the landscape of Kakadu National Park. In: Jones, R. (Ed.), Archaeological Research in Kakadu National Park. Australian National Parks and Wildlife Service and Australian National University Research School of Pacific Studies Joint Publication, Canberra, pp. 229-240.
- Hopf, F.V.L., Colhoun, E.A., Barton, C.E., 2000. Late-glacial and Holocene record of vegetation and climate from Cynthia Bay, Lake St Clair, Tasmania. *Journal of Quaternary Science* **15**, 725-732.
- Hopkins, M.S., Ash, J., Graham, A.W., Head, J., Hewett, R.K., 1993. Charcoal evidence of spatial extent of the eucalyptus woodland expansions and rainforest contractions in North Queensland during the late Pleistocene. *Journal of Biogeography* **20**, 357-372.
- Horne, 2005. Honours Thesis, University of New South Wales, Sydney.
- Horsfall, N., 1987. Living in the rainforest: the prehistoric occupation of north Queensland's humid tropics. PhD Thesis, James Cook University, Townsville.
- Howard, T.M., Hope, G.S., 1970. The present and past occurrence of Beech (Nothofagus cunninghamii Oerst) at Wilsons Promontory, Victoria, Australia. *Proceedings of the Royal Society of Victoria* **83**, 199-210.
- Iles, M.T., 1998. Catchment impacts of mining in the Adelaide River region, Northern Territory, Australia. Honours Thesis, Charles Darwin University, Darwin.
- Ilott, P., 1998. Pollution history of Botany Bay assessed via geochemical analysis of Towra Point sediments. Honours Thesis, University of Wollongong, Wollongong.
- Jenkins, M.A., 1992. A Late Holocene vegetation record from an interdunal swamp, Mornington Peninsula, Victoria. M.Env.Sci. Thesis, Monash University, Melbourne.
- Jenkins, M.A., Kershaw, A.E., 1997. A mid-late Holocene record from an interdunal swamp, Mornington Peninsula, Victoria. *Proceedings of the Royal Society of Victoria* **109**, 133-148.
- Johnson, A.G., 1994. Late Holocene environmental changes on Kurnell Peninsula, NSW. *Proceedings of the Linnean Society of New South Wales* **114**, 119-132.

Johnson, A.G., 1997. PhD Thesis, University of New South Wales, Sydney.

Johnson, A.G., 2000. Fine resolution palaeoecology confirms anthropogenic impact during the Late Holocene in the Lower Hawkesbury Valley, NSW. *Australian Geographer* **31**, 209-235.

Jones, B.G., Killian, H.E., Chenhall, B.E., Sloss, C.R., 2003. Anthropogenic Effects in a Coastal Lagoon: Geochemical Characterization of Burrill Lake, NSW, Australia. *Journal of Coastal Research* **19**, 621-632.

Jordan, G.J., 1997. Evidence of Pleistocene plant extinction and diversity from Regatta Point, western Tasmania, Australia. *Botanical Journal of the Linnean Society* **123**, 45-71.

Jordan, G.J., Carpenter, R.J., Hill, R.S., 1991. Late Pleistocene vegetation and climate near Melaleuca Inlet, south-western Tasmania. *Australian Journal of Botany* **39**, 315-333.

Jordan, G.J., Hill, R.S., 1991. Two new Banksia species from Pleistocene sediments in western Tasmania. *Australian Systematic Botany* **4**, 499-511.

Joyce, E.B., 1989.

Kelly, A., 2006. Reconstructing a vegetation history of the coastal peat plains of Byron Bay using pollen analyses. School of Environmental Science and Management, Southern Cross University, Lismore.

Kemp, J., 1994. Pleistocene-Holocene transition in southern NSW. Honours Thesis, Australian National University, Canberra.

Kenyon, C., 2005. Proceedings of the Royal Society of Victoria 117, 41-59.

Kenyon, C.E., 1989. A Late Pleistocene and Holocene palaeoecological record from Boulder Flat, East Gippsland. Honours Thesis, Monash University, Melbourne.

Kershaw, A.P., 1970. A pollen diagram from Lake Euramoo, north-east Queensland, Australia. *New Phytologist* **69**, 785-805.

Kershaw, A.P., 1971. A pollen diagram from Quincan Crater, North-East Queensland, Australia. *New Phytologist* **70**, 669-681.

Kershaw, A.P., 1974. A long continuous pollen sequence from north-eastern Australia. *Nature* **251**, 222-223.

Kershaw, A.P., 1975. Stratigraphy and pollen analysis of Bromfield Swamp, North Eastern Queensland, Australia. *New Phytologist* **75**, 173-191.

Kershaw, A.P., 1978. Record of last interglacial - glacial cycle from northeastern Queensland. *Nature* **272**, 159-161.

Kershaw, A.P., 1979. Local pollen deposition in aquatic sediments on the Atherton Tableland, North-Eastern Australia. *Australian Journal of Ecology* **4**, 253-263.

Kershaw, A.P., 1983. A Holocene pollen diagram from Lynch's Crater, north-eastern Queensland, Australia. *New Phytologist* **94**, 669-682.

Kershaw, A.P., 1986. Climatic change and aboriginal burning in northeast Australia during the last two glacial interglacial cycles. *Nature* **322**, 47-49.

Kershaw, A.P., 1994. Pleistocene vegetation of the humid tropics of northeastern Queensland, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **109**, 399-412.

Kershaw, A.P., Bulman, D., 1996. A preliminary application of the analogue approach to the interpretation of late Quaternary pollen spectra from southeastern Australia. *Quaternary International* **33**, 61-71.

Kershaw, A.P., D'Costa, D.M., McEwen Mason, J.R.C., Wagstaff, B.E., 1991. Palynological evidence for Quaternary vegetation and environments of mainland southeastern Australia. *Quaternary Science Reviews* **10**, 391-404.

Kershaw, A.P., Dunkerley, D.L., Quinn, M., Southern, W., 1979. A Scientific and environmental investigation of Jackson's Bog, southern tablelands of N.S.W. Unpublished Report to National Parks and Wildlife Service of New South Wales, Melbourne.

Kershaw, A.P., Green, J.E., 1983. Tawonga Bog revisited: the history of a low altitude peat deposit. *Victorian Naturalist* **100**, 256-259.

Kershaw, A.P., Sluiter, I.R., 1982. Late Cenozoic pollen spectra from the Atherton Tableland, north-eastern Australia. *Australian Journal of Botany* **30**, 279-295.

Kershaw, A.P., Strickland, K.M., 1989. The development of alpine vegetation on the Australian mainland. In: Good, R. (Ed.), The scientific significance of the Australian Alps: the proceedings of the first Fenner Conference on the Environment. Australian Alps National Parks Liaison Committee in association with the Australian Academy of Science, Canberra, pp. 113-126.

Kershaw, A.P., Tibby, J.C., Penny, D., Yesdani, H., Walkley, R., Cook, E., Johnston, R., 2004. Latest Pleistocene and Holocene vegetation and environmental history of the Western Plains of Victoria. *Proceedings of the Royal Society of Victoria* **116**, 141-163.

Knight, C., 2006. A Paleolimnological study of Lake Arragan. School of Environmental Science and Management, Southern Cross University, Lismore.

Kodela, P.G., 1990. Modern Pollen Rain From Forest Communities on the Robertson Plateau, New South Wales. *Australian Journal of Botany* **38**, 1-24.

Kodela, P.G., 1992. Rainforest, pollen and palaeoecological studies in the Robertson area. *Eucryphia* **5**, 5-8.

Kodela, P.G., 1996. PhD Thesis, University of New South Wales, Sydney.

Kodela, P.G., Dodson, J.R., 1989. A late Holocene vegetation and fire record from Ku-ring-gai Chase National Park, New South Wales. *Proceedings of the Linnean Society of New South Wales* **110**, 317-326.

Kodela, P.G., Hope, G.S., 1992. Wingecarribee Swamp: Statement of Significance. Report to NSW National Trust, Sydney.

Krull, E.S., Bestland, E.A., Skjemstada, J.O., Parr, J.F., 2006. Geochemistry (δ13C, δ15N, 13C NMR) and residence times (14C and OSL) of soil organic matter from red-brown earths of South Australia: Implications for soil genesis. *Geoderma* **132**, 344-360.

Ladd, P.G., 1971. A Pollen Diagram from Five Mile Beach Swamp, Wilsons Promontory. Honours Thesis, University of Melbourne, Melbourne.

Ladd, P.G., 1976. Past and present vegetation of the Lancefield area, Victoria. *The Artefact* 1, 113-127.

Ladd, P.G., 1978. Vegetation history at Lake Curlip in Lowland Eastern Victoria, from 5200 B.P. to Present. *Australian Journal of Botany* **26**, 393-414.

Ladd, P.G., 1979. A Holocene vegetation record from the eastern side of Wilsons Promontory, Victoria. *New Phytologist* **82**, 265-276.

Ladd, P.G., 1979. Past and Present Vegetation on the Delegate River in the Highlands of Eastern Victoria. I. Present Vegetation. *Australian Journal of Botany* **27**, 167-184.

Ladd, P.G., 1979. Past and Present Vegetation on the Delegate River in the Highlands of Eastern Victoria. II. Vegetation and Climatic History From 12,000 B.P. To Present. *Australian Journal of Botany* 27, 185-202.

Ladd, P.G., Orchista, D.W., Joyce, E.B., 1992. Holocene vegetation history of Flinders Island. *New Phytologist* **122**, 757-767.

Lane, C., 1999. The use of diatoms as biological indicators of water quality in two coastal dune lakes, northern NSW. Honours Thesis, Southern Cross University, Lismore.

Lane, C., 2004. The use of diatoms as biological indicators of water quality, and for environmental reconstruction, in southeast Tasmania, Australia. PhD Thesis, University of Tasmania, Hobart.

Langeveld, M., in prep. Nutrient cycling in a modified tropical macrotidal environment. PhD Thesis, Charles Darwin University, Darwin.

Leahy, P.J., Tibby, J., Kershaw, A.P., Heijnis, H., Kershaw, J.S., 2005. The impact of European settlement on Bolin Billabong, a Yarra River floodplain lake, Melbourne, Australia. *River Research and Applications* **21**, 131-149.

Leech, 2002. Honours Thesis, University of New South Wales, Sydney.

Lewis, T., 2005. A Holocene multi-proxy palaeoecological record from Lake Surprise, Mt Eccles, south-western Victoria. Honours Thesis, Monash University, Melbourne.

Ling, H., Tyler, P., 1995. The Staurastrum Tohopekaligense Species Cluster. *Algological Studies* **76**, 27-60.

Ling, H., Tyler, P., 2000. Australian Freshwater Algae (exclusive of diatoms). Gebrüder Borntraeger, Berlin.

Litchfield, W.H., 1975. Palynological studies in Brigalow (Acacia harpophylla F.Muell) communities: Pollen analysis in Gilgai soil under the primary forest. *Australian Journal of Botany* **23**, 355-371.

Lloyd, P.J., Kershaw, A.P., 1997. Late Quaternary vegetation and early Holocene quantitative climate estimates from Morwell Swamp, Latrobe Valley, southeastern Australia. *Australian Journal of Botany* **45**, 549-563.

Logan, B., 2003. Paleo-environmental reconstruction of a wetland in west Byron Bay, and the establishment of pre-European settlement environmental conditions. Honours Thesis, Southern Cross University, Lismore.

Longmore, M.E., 1986. Modern and ancient sediments - database for management of aquatic ecosystems and their catchments. In: De Deckker, P., Williams, W.D. (Eds.), Limnology in Australia. Dr Junk Publishers, Dordrecht.

Longmore, M.E., 1997. Quaternary Palynological Records from Perched Lake Sediments, Fraser Island, Queensland, Australia: Rainforest, Forest History and Climatic Control. *Australian Journal of Botany* **45**, 507-526.

Longmore, M.E., 1998. The mid-Holocene "Dry" Anomaly on the Mid-Eastern Coast of Australia: Calibration of Palaeowater Depth as a Surrogate for Effective Precipitation using Sedimentary Loss on Ignition in the Perched Lake Sediments of Fraser island, Queensland. *Palaeoclimates - Data and Modelling* **3**, 135-160.

Longmore, M.E., Heijnis, H., 1999. Aridity in Australia: Pleistocene records of paleohydrological and palaeoecological change from the perched lake sediments of Fraser Island, Queensland, Australia. *Quaternary International* **57-58**, 35-47.

Longmore, M.E.M., O'Leary, B.M., Rose, C.W., 1983. Caesium-137 profiles in the sediments of a partical-meromictic lake on Great Sandy Island (Fraser Island), Queensland, Australia. *Hydrobiologia* **103**, 21-27.

Luly, J., 1990. Holocene palaeoenvironments at Lake Tyrrell, Victoria. PhD Thesis, Australian National University, Canberra.

Luly, J.G., 1993. Holocene Palaeoenvironments near Lake Tyrell, semi-arid northwestern Victoria, Australia. *Journal of Biogeography* **20**, 587-598.

Luly, J.G., 1995. Holocene Palaeoenvironments at Lake Tyrrell: Response to Sluiter and Parsons. *Journal of Biogeography* **22**, 152-156.

MacGregor, A.J., Gell, P.A., Wallbrink, P.J., Hancock, G., 2005. Natural and post-disturbance variability in water quality of the lower Snowy River floodplain, Eastern Victoria, Australia. River *Research and Applications* **21**, 201-213.

Macphail, M., Hope, G., Anderson, A., 2001. Polynesian Plant Introductions in the southwest Pacific: Initial Pollen Evidence for Norfolk Island. *Records of the Australian Museum Supplement* **27**, 123-134.

Macphail, M., Neale, J., 1996. Kingston Swamp, Norfolk Island: a unique natural archive of SW Pacific Prehistory? Interim Report prepared for the Australian Heritage Commission, Canberra.

Macphail, M.K., Unpublished report.

Macphail, M.K., 1975. Late Pleistocene Environments in Tasmania. Search 6, 295-300.

Macphail, M.K., 1979. Vegetation and climates in southern Tasmania since the last glaciation. *Quaternary Research* **11**, 306-341.

Macphail, M.K., 1986. Over the Top - pollen based reconstructions of past alpine floras and vegetation in Tasmania. In: Barlow, B.A. (Ed.), Flora and Fauna of Alpine Australasia: ages and origins. CSIRO in association with Australian Systematic Botany Society, Melbourne pp. 173-203.

Macphail, M.K., 1999. A hidden cultural landscape: colonial Sydney's plant microfossil record. *Australian Historical Archaeology* **17**, 79-109.

Macphail, M.K., Higginbotham, E., Davey, B.G., 1988. An early colonial garden landscape at Sydney Cove; pollen and soil. *Search* **19**, 141-144.

Macphail, M.K., Hope, G.S., 1997. Palynological report, Core Site KCA, Kingston Swamp, Norfolk Island. Unpublished report to Australian Heritage Commission, Canberra.

Magee, J.W., Bowler, J.M., Miller, G.H., Williams, D.L.G., 1995. Stratigraphy, Sedimentology, Chronology and Paleohydrology of Quaternary Lacustrine Deposits at Madigan Gulf, Lake Eyre, South Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **113**, 3-42.

Markgraf, V., Bradbury, J.P., Busby, J.R., 1986. Paleoclimates in Southwestern Tasmania during the last 13,000 years. *Palaios* 1, 368-380.

Martin, A.R.H., 1972. The Depositional environment of the organic deposits on the foreshore at North Dee Why, New South Wales. *Proceedings of the Linnean Society of New South Wales* **96**, 278-281.

Martin, A.R.H., 1986. Late Glacial and early Holocene vegetation of the alpine zone, Kosciuszko National Park. In: Barlow, B.A. (Ed.), Flora and Fauna of Alpine Australia: ages and origins. CSIRO, Melbourne, pp. 161-170.

Martin, A.R.H., 1986. Late glacial and Holocene alpine pollen diagrams from the Kosciuszko National Park, New South Wales, Australia. *Review of Palaeobotany and Palynology* **47**, 367-409.

Martin, A.R.H., 1986. Palaeoecology, Palaeolimnology, Palynology and Coastal Environment - areas of neglect in coastal studies? In: Frankel, E., Keene, J.B., Waltho, A.E. (Eds.), Recent Sediments in eastern Australia; Marine through Terrestrial. Geological Society of Australia, Sydney.

Martin, A.R.H., 1994. Kurnell Fen: an eastern Australian coastal wetland, its Holocene vegetation, relevant to sea-level change and aboriginal land use. *Review of Palaeobotany and Palynology* **80**, 311-332.

Martin, H.A., McMinn, A., 1994. Late Cainozoic vegetation history of north-western Australia, from the palynology of a deep sea Core (ODP 765). *Australian Journal of Botany* **42**, 95-102.

Mason, 2004. Honours Thesis, University of New South Wales, Sydney.

Maxwell, C., Tyler, P., 2006. Recolonisation of Lagoon of Islands, Tasmania, by Baumea Arthrophylla: The First Step in Regeneration of a Unique Ecosystem? *Proceedings of the Royal Society of Tasmania* **140**, 31-34.

McCarthy, L., Head, L., 2001. Holocene variability in semi-arid vegetation: new evidence from Leporillus middens from the Flinders Ranges, South Australia. *The Holocene* **11**, 681-689.

McCarthy, L., Head, L., Quade, J., 1996. Holocene palaeoecology of the northern Flinders ranges, South Australia, based on stick-nest rat (Leporillus spp.) middens: a preliminary overview. *Palaeogeography, Palaeoclimatology, Palaeoecology* **123**, 205-218.

McConnell, K., 1998. The prehistoric use of Chenopodiaceae in Australia: evidence from Carpenter's Gap shelter 1 in the Kimberley, Australia. *Vegetation History and Archaeobotany* **7**, 179-188.

McConnell, K., O'Connor, S., 1997. 40,000 year record of food plants in the southern Kimberley Ranges, Western Australia. *Australian Archaeology* **45**, 20-31.

McGrath, R.J., Boyd, W.E., 1998. Holocene vegetation history of Bundjalung National Park and Bungawalbin Creek, north-eastern New South Wales. *Australian Geographer* **29**, 205-221.

McKenzie, G.M., 1989. Late Quaternary vegetation and climate in the central Highlands of Victoria, with special reference to Nothofagus cunninghamii (Hook.) Oerst. rainforest. PhD Thesis, Monash University, Melbourne.

McKenzie, G.M., 1997. The late Quaternary vegetation history of the south-central highlands of Victoria, Australia:1. Sites above 900m. *Australian Journal of Ecology* **22**, 19-36.

McKenzie, G.M., 2002. The late Quaternary vegetation history of the south-central highlands of Victoria, Australia. II. Sites below 900 m. *Austral Ecology* **27**, 32-54.

McKenzie, G.M., Busby, J.R., 1992. A Quantitative estimate of Holocene climate using a bioclimatic profile of Nothofagus cunninghamii (Hook.) Oerst. *Journal of Biogeography* **19**, 531-540.

McKenzie, G.M., Kershaw, A.P., 1997. A Vegetation History and Quantitative Estimate of Holocene Climate from Chapple Vale, in the Otway Region of Victoria, Australia. *Australian Journal of Botany* **45**, 565-581.

McKenzie, G.M., Kershaw, A.P., 2000. The last glacial cycle from Wyelangta, the Otway region of Victoria, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **155**, 177-193.

McMinn, A., Augustinus, P., Barton, C., 2003. Diatom analysis of late Holocene sediment cores from Macquarie Harbour, Tasmania. *Alcheringa* **27**, 135-153.

McMinn, A., Hallegraeff, G., Roberts, J., Smith, J., Lovell, A., Jenkinson, A., Heijnis, H., 2001. Recent introduction of Gymnodinium catenatum to Port Lincoln, South Australia. *Harmful Algal Blooms* **2000**, 477-480.

McMinn, A., Heijnis, H., Murray, A., Hallegraeff, G., 2004. Diatom and dinoflagellate assemblages of the Hawkesbury River, NSW, over the last two centuries: evidence for changes in hydrology. *Alcheringa* **28**, 505-514.

McNess, A., 1988. A fine resolution vegetation history from a Melbourne metropolitan park. Honours Thesis, Monash University, Melbourne.

Metcalfe, G.D., 1999. Without a paddle: A study of the effects of land use and urban development in the Erskine Creek catchment. Honours Thesis, University of Wollongong, Wollongong.

Mooney, S., 1997. A fine-resolution palaeoclimatic reconstruction of the last 2000 years from Lake Keilambete, south-eastern Australia. *The Holocene* **7**, 139-149.

Mooney, S.D., Maltby, E.L., 2006. Two proxy records revealing the late Holocene fire history at a site on the central coast of New South Wales, Australia. *Austral Ecology* **31**, 682-695.

Mooney, S.D., Radford, K.L., Hancock, G.J., 2001. Clues to the 'burning question': pre-European fire in the Sydney coastal region from sedimentary charcoal and palynology. *Ecological Management and Restoration* **2**, 203-212.

Mooney, S.D., Watson, J.R., Dodson, J.R., 1997. Late Holocene environmental change in an upper montane area of the Snowy Mountains, New South Wales. *Australian Geographer* **28**, 185-200.

Mooney, S.D., Webb, M., Attenbrow, V., 2007. A Comparison of Charcoal and Archaeological Information to Address the Influences on Holocene Fire Activity in the Sydney Basin. *Australian Geographer* **38**, 177-194.

Morris, A., 1996. The Biogeography of Plant Communities at Woodman Point, WA. Postgraduate Diploma of Science Thesis, University of Western Australia, Perth.

Moss, P., 1994. Late Holocene environments of Den Plain, north-western Tasmania. Honours Thesis, University of Melbourne, Melbourne.

Moss, P.T., Kershaw, A.P., 2000. The last glacial cycle from the humid tropics of northeastern Australia: comparison of a terrestrial and marine record. *Palaeogeography, Palaeoclimatology, Palaeoecology* **155**, 155-176.

Napoli, M., 1996. The contaminant chronologies of Prospect Creek and Salt Pan Creek as recorded in their sediments. Honours Thesis, University of Sydney.

Newsome, J.C., 1999. Pollen-vegetation relationships in semi-arid southwestern Australia. *Review of Palaeobotany and Palynology* **106**, 103-119.

Newsome, J.C., Pickett, E.J., 1993. Palynology and palaeoclimatic implications of two Holocene sequences from southwestern Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **101**, 245-261.

Noakes, A.J., 1998. A study of the land use change and fire record of the Thirlmere Lakes area since European settlement as provded by the sediment record. Honours Thesis, University of Wollongong, Wollongong.

Noble, M.G., 1981. Vegetation organisation along altitudinal gradients in Tasmania. PhD Thesis, Australian National University, Canberra.

Oakes, G.M., 1981. The Jackson's Bog peat deposits. Unpublished New South Wales Geological Survey Report GS 1981/596, Sydney.

O'Connor, S., 1992. The timing and nature of prehistoric island use in northern Australia. *Archaeology in Oceania* **27**, 49-60.

Ogden, R.W., 1996. The impacts of farming and river regulation on billabongs of the southeast Murray Basin, Australia. PhD Thesis, Australian National University, Canberra.

Ogden, R.W., 1997. The effects of European settlement on the biodiversity of chydorid Cladocera in billabongs of the south-east Murray Basin. *Memoirs of the Museum of Victoria* **56**, 505-511.

Ogden, R.W., 2000. Modern and historical variation in aquatic macrophyte cover of billabongs associated with catchment development. *Regulated Rivers: Research and Management* **16**, 497-451.

Organo, R.J., 2000. The spatial and temporal distribution of pollution in the Cox's River, NSW. Honours Thesis, University of Wollongong, Wollongong.

- Packwood, A., 1999. An environmental investigation of Lake Conjola, South Coast, NSW. Honours Thesis, University of Wollongong, Wollongong.
- Panayotou, K., 2004. Geomorphology of the Minnamurra River estuary, southeastern Australia: evolution and management of a barrier estuary. PhD Thesis, University of Wollongong, Wollongong.
- Parr, J.F., 2001. Phytolith and starch analysis of sediment samples from two archaeological sites on Dauar Island, Torres Strait. Southern Cross University, Lismore.
- Parr, J.F., 2002. A Pre-Lapita Landscape at Numundo, West New Britain, Papua New Guinea as Indicated by Fossil Phytoliths, The 17th Congress of the Indo-Pacific Prehistory Association. Academia Sinica, Taipei, Taiwan.
- Parr, J.F., 2002. West Byron Wetlands Holocene Vegetation History Report. Southern Cross University, Lismore.
- Parr, J.F., 2003. A study of Palaeo-Landscapes in the Numundo region of West New Britain, Papua New Guinea, as indicated by Fossil Phytolith Analysis. PhD Thesis, Southern Cross University, Lismore.
- Parr, J.F., 2006. Phytolith analysis of samples from Swears Island, Gulf of Carpentaria, Queensland. Southern Cross University, Lismore.
- Parr, J.F., Carter, M., 2003. Phytolith and starch analysis of sediment samples from two archaeological sites on Dauar Island, Torres Strait. *Vegetation History and Archaeobotany* **12**, 131-141.
- Parr, J.F., Kerr, G., Arthur, J., Taffs, K.H., 2003. Fires and their implications for acidic peatlands, northeastern NSW, Ecological Society of Australia Conference. Ecological Society of Australia, Armidale.
- Parr, J.F., Kerr, G., Arthur, J., Taffs, K.H., in press. Fires and their implications for the acidic peatlands of northeastern NSW. In: Korstanje, A., Babot, P. (Eds.). Altamira and British Archaeological Series.
- Parr, J.F., Lentfer, C.J., Boyd, W.E., 2001. Spatial analysis of phytolith assemblages at an archaeological site in West New Britain, Papua New Guinea. In: Clark, G.R., Anderson, A.J., Vunidilo., T. (Eds.), The Archaeology of Lapita Dispersal in Oceania. Pandanus Books, Australian National University, pp. 125-134.
- Parr, J.F., Lentfer, C.J., Boyd, W.E., 2001. Spatial patterning of a Lapita landscape at an archaeological site in the West New Britain Province, Papua New Guinea. *Domodomo (Journal of the Fiji Museum)* **13**, 7-18.
- Parr, J.F., Sullivan, L.A., 2005. Soil carbon sequestration in Phytoliths. *Soil Biology and Biochemistry* **37**, 117-124.

Paterson, C.G., Walker, K.F., 1974. Recent history of Tanytarsus barbitarsus Freeman (Diptera: Chironomidae) in the sediments of a shallow saline lake. *Marine and Freshwater Research* **25**, 315-325.

Pearson, S., 1999. Late Holocene biological records from the middens of stick-nest rats in the central Australian arid zone. *Quaternary International* **59**, 39-46.

Penny, D., 1993. Palaeoecology and human impact at Lake Coleman, East Gippsland, Victoria. Honours Thesis, Monash University, Melbourne.

Peters, G.M., Maher, W.A., Jolley, D., Carroll, B.I., Gomes, V.G., Jenkinson, A.V., McOrist, G.D., 1999. Selenium contamination, redistribution and remobilisation in sediments of Lake Macquarie, NSW. *Organic Geochemistry* **30**, 1287-1300.

Pfitzner, J., Brunskill, G., Zagorskis, I., 2004. 137Cs and excess 210Pb deposition patterns in estuarine and marine sediment in the central region of the Great Barrier Reef Lagoon, north-eastern Australia. *Journal of Environmental Radioactivity* **76**, 81-102.

Pickett, E.J., Harrison, S.P., Hope, G., Harle, K., Dodson, J.R., Kershaw, A.P., Prentice, I.C., Backhouse, J., Colhoun, E.A., D'Costa, D., Flenley, J., Grindrod, J., Haberle, S., Hassell, C., Kenyon, C., Macphail, M., Martin, H., Martin, A.H., McKenzie, M., Newsome, J.C., Penny, D., Powell, J., Raine, J.I., Southern, W., Stevenson, J., Sutra, J.-P., Thomas, I., van der Kaars, S., Ward, J., 2004. Pollen-based reconstructions of biome distributions for Australia, Southeast Asia and the Pacific (SEAPAC region) at 0, 6000 and 18,000 14C yr BP. *Journal of Biogeography* 31, 1381-1444.

Pickett, E.J., Newsome, J.C., 1997. Eucalyptus (Myrtaceae) pollen and its potential role in investigations of Holocene environments in Southwestern Australia. *Review of Palaeobotany and Palynology* **98**, 187-205.

Pittock, 1989. Honours Thesis, Monash University, Melbourne.

Pola, J., 1993. Late Pleistocene pollen and vegetation history from Lake Selina, western Tasmania. Honours Thesis, University of Newcastle, Newcastle.

Polach, H., Singh, G., 1980. Contemporary 14C levels and their significance to the sedimentary history of Bega Swamp, North South Wales. *Radiocarbon* **22**, 398-409.

Prebble, M., Sim, R., Finn, J., Fink, D., 2005. A Holocene pollen and diatom record from Vanderlin Island, Gulf of Carpentaria, lowland tropical Australia. *Quaternary Research* **64**, 357-371.

Radclyffe, E., 1993. Fire and people in Jervis Bay: toward a fire history in an occupied landscape. Honours Thesis, Australian National University, Canberra.

Radke, L.C., Prosser, I., Robb, M., Brooke, B., Fredericks, D., Douglas, G.B., Skemstad, J., 2004. The relationship between sediment and water quality, and riverine sediment loads in the wave-

dominated estuaries of south-west Western Australia. *Marine and Freshwater Research* **55**, 581–596.

Rainbow, C.M., 1999. Chronology of sediments and spatial pollutant distribution patterns in the Werri Berri Creek catchment, Wollondilly Shire, NSW. Honours Thesis, University of Wollongong, Wollongong.

Raine, J.I., 1974. Pollen Sedimentation in Relation to Quaternary Vegetation History of the Snowy Mountains of New South Wales. PhD Thesis, Australian National University, Canberra.

Raine, J.I., 1982. Dimictic thermal regime and morphology of Blue Lake in the Snowy Mountains of New South Wales. *Australian Journal of Marine and Freshwater Research* **33**, 1119 - 1122.

Ralph, T.J., 2001. The rate and distribution of sedimentation adjacent to a distributary channel in the Macquarie Marshes and the implications for processes of channel avulsion. Honours Thesis, Macquarie University, Sydney.

Ralph, T.J., in prep. Channel breakdown and floodplain wetland morphodynamics in the Macquarie Marshes. PhD Thesis, Macquarie University, Sydney.

Reid, M., 1989. Palaeoecological Changes at Lake Wellington, Gippsland Lakes, Victoria, during the late Holocene: A Study of the Development of a Coastal Lake Ecoystem. Honours Thesis, Monash University, Melbourne.

Reid, M., 1998. A fine resolution palaeoecology of recent sediments from Lake Wellington, Gippsland Lakes, Victoria. In: Newall, P. (Ed.), Proceedings of the First Australian Diatom Workshop. Deakin University School of Ecology and Environment, Warrnambool.

Reid, M., 2002. A diatom-based palaeoecological study of two billabongs on the Goulburn River floodplain, southeast Australia. In: John, J. (Ed.), Proceedings of the 15th International Diatom Symposium. Königstein, Koeltz Scientific Books, Perth.

Reid, M., Fluin, J., Ogden, R., Tibby, J., Kershaw, P., 2002. Long-term perspectives on human impacts on floodplain-river ecosystems, Murray-Darling Basin, Australia. *Verhandlungen der internationalen Vereinigung für theoretische und angewandte Limnologie* **28**, 1-7.

Reid, M., Sayer, C.D., Kershaw, A.P., Heijnis, H., in press. Palaeolimnological evidence for submerged plant loss in a floodplain lake associated with accelerated catchment soil erosion (Murray River, Australia). *Journal of Paleolimnology*.

Reid, M.A., Tibby, J.C., Penny, D., Gell, P.A., 1995. The use of diatoms to assess past and present water quality. *Australian Journal of Ecology* 20, 57-64.

Roach, A., 2005. A Palaeoecological Study Covering the Past 30,000 years of Tyrendarra Swamp, Western Plains, Victoria. Honours Thesis, Monash University, Melbourne.

Robbie, 1998. Honours Thesis, University of New South Wales, Sydney.

Robertson, M., 1986. Fire Regimes and Vegetation Dynamics: A Case Study Examining the Potential of the Fire Resolution Palaeoecological Techniques. Honours Thesis, Monash University, Melbourne.

Rogers, K., Saintilan, N., Heijnis, H., 2005. Mangrove Encroachment of Salt Marsh in Western Port Bay, Victoria: the Role of Sedimentation, Subsidence, and Sea Level Rise. *Estuaries* **28**, 551-559.

Rolfe, S., Kew, P., Tyler, P., 2001. Reconnaissance Limnology of Tasmania VI. Biogeographical Significance of Physicochemical Features of Coastal Lagoons of the Islands of Bass Strait. *Archiv für Hydrobiologie* **150**, 693-704.

Rowe, C., 2007. A palynological investigation of Holocene vegetation change in Torres Strait, seasonal tropics of northern Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **251**, 83-103.

Rowe, R.K., 1968. Rare bog community at Tawonga, Victoria. Victorian Naturalist 85, 123-124.

Rowell, M.V., Jordan, G.J., Barnes, R.W., 2001. An in situ, late Pleistocene Melaleuca fossil forest at Coal Head, western Tasmania, Australia. *Australian Journal of Botany* **49**, 235-244.

Russell-Smith, J., 1985. A record of change: studies of Holocene vegetation history in the South Alligator River region, Northern Territory. *Proceedings of the Ecological Society of Australia* 13, 191-202.

Sabbe, K., Vanhoutte, K., Lowe, R.L., Biggs, B.J.F., Bergey, E., Francoeur, S., Hodgson, D., Vyverman, W., 2001. Six new Actinella (Bacillariophyta) species from Papua New Guinea, Australia and New Zealand: further evidence for widespread diatom endemism in the Australasian region. *European Journal of Phycology* **36**, 321-340.

Sabbe, K., Vanhoutte, K., Vyverman, W., 2000. Actinella comperei sp. nov. (Eunotiaceae, Bacillariophyta): a new endemic freshwater diatom from Tasmania (Australia). *Systematics and Geography of Plants* **70**, 237-243.

Salas, M.R., 1981. A Reconnaissance Survey of the Vegetation and Sedimentary Record of Lake Tali Karng. Honours Thesis, Monash University, Melbourne.

Salas, M.R., 1983. Long distance pollen transport over the southern Tasman Sea: evidence from Macquarie Island. *New Zealand Journal of Botany* **21**, 285-292.

Salas, M.R., Kershaw, A.P., 1982. Pollen analysis of sediment samples from Blue Lake, South Australia. CSIRO Division of Soils, Adelaide.

Saunders, K.M., 2005. Assessing human impacts on southeast Australian coastal lagoons: application of a diatom-nutrient transfer function. Honours Thesis, University of Tasmania, Hobart.

- Saunders, K.M., Hodgson, D.A., Harrison, J., McMinn, A., in press. Palaeoecological tools for improving the management of coastal ecosystems: a case study from Lake King (Gippsland Lakes) Australia. *Journal of Paleolimnology*.
- Saunders, K.M., McMinn, A., Roberts, D., Hodgson, D.A., Heijnis, H., 2007. Recent human-induced salinity changes in Ramsar-listed Orielton Lagoon, south-east Tasmania, Australia: a new approach for coastal lagoon conservation and management. *Aquatic Conservation: Marine and Freshwater Ecosystems* **17**, 51-70.
- Scarlett, N.H., Hope, G.S., Calder, D.M., 1974. Natural History of the Hogan Group. 3. Floristics and Plant Communities. *Papers and Proceedings of the Royal Society of Tasmania* **107**, 83-98.
- Seen, A., Townsend, A., Atkinson, B., Ellison, J., Harrison, J., Heijnis, H., 2004. Determining the history and sources of contaminants in sediments in the Tamar Estuary, Tasmania, Australia, using 210Pb dating and stable Pb isotopes. *Environmental Chemistry* 1, 49-54.
- Selkirk, D.R., Selkirk, P.M., Griffin, K., 1982. Palynological Evidence for Holocene Environmental Changes and Uplift on Wireless Hill, Macquarie Island. *Proceedings of the Linnean Society of New South Wales* **107**, 1-17.
- Shulmeister, J., 2: 107-116., 1992. A Holocene pollen record from lowland tropical Australia. *The Holocene* **2**, 107-116.
- Sim, A.K., 2004. The investigation of vegetation thickening in central Queensland using palaeoecological techniques. Honours Thesis, University of New South Wales, Sydney.
- Singh, G., 1983. Late Quaternary vegetation and lake level record from Lake George, New South Wales: 18±2Ka, 15-10 Ka. In: Chappell, J.M.A., Grindrod, A. (Eds.), Proceedings of the First CLIMANZ Conference, Howman's Gap, 1981. Department of Biogeography and Geomorphology, Research School of Pacific Studies, Australian National University, Canberra.
- Singh, G., Geissler, E.A., 1985. Late Cainozoic history of vegetation, fire, lake levels and climate at Lake George, New South Wales, Australia. *Philosophical Transactions of the Royal Society of London B* **311**, 379-447.
- Singh, G., Kershaw, A.P., Clark, R., 1981. Quaternary vegetation and fire history in Australia. In: Gill, A.M., Groves, R.A., Noble, I.R. (Eds.), Fire and the Australian Biota. Australian Academy of Science, Canberra, pp. 23-54.
- Singh, G., Luly, J., 1991. Changes in vegetation and seasonal climate since the last full glacial at Lake Frome, South Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **84**, 75-86.
- Sloss, C., 2001. Holocene stratigraphic evolution and recent sedimentological trends, Lake Illawarra, NSW. Honours Thesis, University of Wollongong, Wollongong.
- Sluiter, I.R.K., Parsons, R.F., 1995. On the Holocene palaeoenvironmental record from Lake Tyrrell, northwestern Victoria, Australia: A reply. *Journal of Biogeography* **22**, 149-156.

Smeulders, D., 1999. Honours Thesis, University of New South Wales, Sydney.

Smith, J., Hamilton, T., 1985. Modelling of 210Pb behaviour in the catchment and sediment of Lake Tali Karng, Victoria, and estimation of recent sedimentation rates. *Australian Journal of Marine and Freshwater Research* **36**, 15-22.

Smith, J.D., 1996. Study of sedimentation rates in Port Phillip Bay using lead-210. CSIRO Environmental Projects Office, Canberra.

Smith, M., 1982. Late Pleistocene Zamia Exploitation in southern Western Australia. *Archaeology in Oceania* **17**, 117-121.

Smith, M., 1996. Revisiting Pleistocene Macrozamia. Australian Archaeology 42, 52-53.

Southern, W., 1982. Late Quaternary vegetation and environments of Jackon's Bog and the Monaro Tablelands, New South Wales. MSc Thesis, Monash University, Melbourne.

Strickland, K.M., 1980. Studies on the relationship between the pollen rain and the vegetation in the vicinity of Cascade Creek on the Baw Plateau, Victoria. Honours Thesis, Monash University, Melbourne.

Sweller, S., Martin, H.A., 1996. History of the vegetation at Burraga Swamp, Barrington Tops National Park, Upper Hunter River Region, NSW. *Proceedings of the Linnean Society of New South Wales* **118**, 23-50.

Sweller, S., Martin, H.A., 2001. A 40,000 year vegetation history and climatic interpretations of Burraga Swamp, Barrington Tops, New South Wales. *Quaternary International* **83-85**, 233-244.

Taffs, K.H., 2001. Diatoms as indicators of wetland salinity in the Upper South East of South Australia. *The Holocene* **11**, 281-290.

Taffs, K.H., Farago, L.J., Heijnis, H., Jacobsen, G., in press. A diatom based Holocene record of human impact from a coastal environment: Tuckean Swamp, eastern Australia. *Journal of Paleolimnology*.

Taffs, K.H., Parr, J.F., Bolton, K., 2006. Using palaeobotanical techniques to guide peatland restoration. A case study from Byron Bay, Australia. *Ecological Management and Restoration* 7, 132-135.

Thom, B.G., Hails, J.H., Martin, A.R.H., 1969. Radiocarbon evidence against higher postglacial sealevels in eastern Australia. *Marine Geology* **7**, 161-168.

Thomas, A., 1991. Giandarra Bog, NSW - A Natural Record of Environmental Change. Honours Thesis, UNSW@ADFA, Canberra.

Thomas, I., 1992. The Holocene Palaeoecology and archaeology of northeastern Tasmania, Australia. PhD Thesis, University of Tasmania, Hobart.

- Thomas, I., 1995. In: Dixon, G., Aitken, D. (Eds.), 1993 Institute of Australian Geographers Conference. Monash Publications in Geography.
- Thomas, I., 1996. Environmental Changes in Northeastern Tasmania during the Holocene.
- Thoms, M.C., Ogden, R., Reid, M.A., 1999. Establishing the condition of lowland floodplain rivers: a palaeo-ecological approach. *Freshwater Biology* **41**, 407-423.
- Tibby, J., 1996. A mid-to late-Holocene diatom and pollen palaeoecology of the Tuross Lake system, South Coast, New South Wales. Monash Publications in Geography 46, 1-62.
- Tibby, J., 1998. A diatom-based history of Burrinjuck Reservoir (southern New South Wales) and its catchment. In: Newall, P. (Ed.), First Australian Diatom Workshop. Deakin University School of Ecology and Environment, Warrnambool.
- Tibby, J., 1998. A mid-Holocene to present palaeoecological study of the Tuross Lakes system, south coast, New South Wales. In: Newall, P. (Ed.), First Australian Diatom Workshop. Deakin University School of Ecology and Environment, Warrnambool.
- Tibby, J., Gell, P., Hancock, G., Clark, M., submitted. Core correlation and sediment chronology from Kangaroo Creek Reservoir, South Australia: implications for palaeolimnology and catchment management. *Journal of Paleolimnology*.
- Tibby, J., Gell, P.A., Cox, A., Hancock, G., in prep. Water quality change in Upper River Torrens water storages, South Australia: inter-basin transfer as a pollution source. *Freshwater Biology*.
- Tibby, J., Haberle, S.G., 2007. A late glacial to present diatom record from Lake Euramoo, wet tropics of Queensland, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **251**, 46-56.
- Tibby, J., Kershaw, A.P., Builth, H., Philibert, A., White, C., 2006. Environmental Change and Variability in Southwestern Victoria: Changing Constraints and Opportunities for Occupation and Land Use. In: David, B., Barker, B., McNiven, I.J. (Eds.), The Social Archaeology of Australian Indigenous Societies. Aboriginal Studies Press, Canberra, pp. 254-269.
- Tibby, J., Lane, M., Gell, P.A., submitted. Local knowledge as a basis for environmental management: a cautionary tale from Lake Ainsworth, northern New South Wales. *Environmental Conservation*.
- Tibby, J., Reid, M., Fluin, J., Hart, B.T., Kershaw, A.P., 2003. Assessing long-term pH change in an Australian river catchment using monitoring and palaeolimnological data. *Environmental Science and Technology* **37**, 3250-3255.
- Tibby, J.C., 2001. Diatoms as indicators of sedimentary processes in Burrinjuck reservoir, New South Wales, Australia. *Quaternary International* **83-85**, 245-256.
- Tibby, J.C., 2004. Assessing the impact of early colonial Australia on the physical environment: a comment on Gale and Haworth (2002). *Archaeology in Oceania* **39**, 144-148.

- Torgersen, T., Longmore, M.E., 1984. Caesium 137 diffusion in the highly organic sediment of Hidden Lake, Fraser Island, Queensland. *Marine and Freshwater Research* **35**, 537-548.
- Torrence, R., Neall, V.E., Doelman, T., Rhodes, E., McKee, C., Davies, H., Bonetti, R., Gugliemetti, A., Manzoni, A., Oddone, M., Parr, J., Wallace, C., 2004. Pleistocene colonisation of the Bismarck Archipelago: new evidence from West New Britain. *Archaeology in Oceania* **39**, 101-130.
- Treague, J., 2003. Geological context of peat deposition in Cumebin Swamp, NSW. A geological framework for carbon sequestration sudies. School of Environmental Science and Management, Southern Cross University, Lismore.
- Turney, C.S.M., Kershaw, A.P., Moss, P., Bird, M.I., Fifield, L.K., Cresswell, R.G., Santos, G.M., Di Tada, M.L., Hausladen, P.A., Zhou, Y., 2001. Redating the onset of burning at Lynch's Crater (North Queensland): implications for human settlement in Australia. *Journal of Quaternary Science* **16**, 767-771.
- Tyler, P., 1996. Endemism in freshwater algae with specified reference to the Australian region. In: Kristiansen, J. (Ed.), Biogeography of Freshwater Algae. Kluwer Academic Publishers, Belgium, pp. 127-135.
- Tyler, P., 1996. The significance of northeast Tasmania for the biogeography of endemic Australian freshwater algae, Geography of Northeast Tasmania: Records of the Queen Victoria Museum. Queen Victoria Museum and Art Gallery, Tasmania, pp. 133-135.
- Tyler, P., 1996. Three Thousand Years of Running, Jumping and Standing Still. In: Mitchell, B., Day, D. (Eds.), Sustainable Management of Australia's Inland Waters. Deakin University, Melbourne, pp. 67-80.
- Tyler, P., 2001. Lake Pedder A Limnologist's Lifetime View, Proceedings of a Symposium: Lake Pedder Values and Restoration. University of Tasmania, Tasmania.
- Tyler, P., Hodgson, D., Wright, S., Davies, N., 1998. Analysis of fossil pigments from algae and bacteria in meromictic Lake Fidler, Tasmania, and its application to lake management. *Journal of Paleolimnology* **19**, 1-22.
- Tyler, P., Sherwood, J., Magilton, C., Hodgson, D., 1996. Limnological and Geomorphological Considerations Underlying Pedder 2000 the Campaign to Restore Lake Pedder. *Archiv für Hydrobiologie* **136**, 343-361.
- Tyler, P., Terry, C., Howland, M., 2001. Gordon River Meromictic Lakes Assessment: Basslink Integrated Impact Assessment Statement Potential Effects of Changes to Hydro Power Generation. Hydro Tasmania, Australia.
- Tyler, P., Vyverman, W., 1995. The Microbial Marketplace tradeoffs at the chemocline of meriomictic lakes. In: Round, F.E., Chapman, D.J. (Eds.), Progress in Phycological Research. Biopress Ltd, pp. 325-370.

van de Geer, G., Fitzsimons, S.J., Colhoun, E.A., 1989. Holocene to Middle Last Glaciation vegetation history at Newall Creek, western Tasmania. *New Phytologist* **111**, 549-558.

van der Kaars, S., De Deckker, P., 2002. A Late Quaternary pollen record from deep-sea core Fr10/95, GC17 offshore Cape Range Peninsula, northwestern Western Australia. *Review of Palaeobotany and Palynology* **120**, 17-39.

van der Kaars, S., Wang, X., Kershaw, P., Guichard, F., Setiabudi, D.A., 2000. A late Quaternary palaeoecological record from the Banda Sea, Indonesia: patterns of vegetation, climate and biomass burning in Indonesia and northern Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **155**, 135-153.

Van der Kaars, W.A., 1989. Late Quaternary vegetation and climate of Australasia as reflected by the palynology of eastern Indonesian deep sea piston-cores. PhD Thesis, University of Amsterdam, Amsterdam.

Vanhoutte, K., Verleyen, E., Sabbe, K., Kilroy, C., Sterken, M., Vyverman, W., 2006. Congruence and disparity in benthic diatom community structure of small lakes in New Zealand and Tasmania. *Marine and Freshwater Research* **57**, 789-801.

Vanhoutte, K., Vyverman, W., 2002. Een paleo-ecologische studie van Lake Dobson en Lake Perry (Tasmanië, Australië). *Diatomededelingen* **26**.

Vyverman, W., 1996. The Indo-Malaysian North-Australian phycogeographical region revised. In: Kristiansen, J. (Ed.), Biogeography of Freshwater Algae. Kluwer Academic Publishers, Dordrecht and Boston, pp. 107-120.

Vyverman, W., Muylaert, K., Verleyen, E., Sabbe, K., 2007. Ecology of non-marine microalgae - Lakes and large rivers. In: McCarthy, P.M., Orchard, A.E. (Eds.), The Algae of Australia (Introduction). CSIRO Publishing, Melbourne, pp. 459-475.

Vyverman, W., Sabbe, K., Mann, D., Vyverman, R., Hodgson, D., Vanhoutte, K., Kilroy, C., 1998. Eunophora gen. nov. (Bacillariophyta), a new genus of amphoroid diatoms from Tasmanian highland lakes with close affinities to the genus Eunotia. *European Journal of Phycology* **33**, 95-111.

Vyverman, W., Sabbe, K., Vyverman, R., 1997. Five new freshwater species of Biremis (Bacillariophyta) from Tasmania. *Phycologia* **36**, 91-102.

Vyverman, W., Verleyen, E., Sabbe, K., 2007. Diversity and Biogeography of Freshwater Microalgae. In: McCarthy, P.M., Orchard, A.E. (Eds.), The Algae of Australia (Introduction). CSIRO Publishing, Melbourne, pp. 580-593.

Vyverman, W., Vyverman, R., Hodgson, D., P.A., T., 1995. Diatoms from Tasmanian mountain lakes: A reference data-set (TASDIAT) for environmental reconstruction and a systematic and autecological study. *Bibliotheca Diatomologica* **33**, 1-193.

- Vyverman, W., Vyverman, R., Tyler, P.A., Rajendran, V.S., 1996. Distribution of benthic diatom assemblages in Tasmanian highland lakes and their possible use as indicators of environmental changes. *Canadian Journal of Fisheries and Aquatic Sciences* **53**, 493-508.
- Walker, D., 1999. Some physical and chemical features of two upland crater lakes in tropical north-eastern Australia. *Marine and Freshwater Research* **50**, 159-177.
- Walker, D., 2000. Pollen input to, and incorporation in, two crater lakes in tropical northeast Australia. *Review of Palaeobotany and Palynology* **111**, 253-283.
- Walker, D., 2007. Holocene sediments of Lake Barrine, north-east Australia, and their implications for the history of lake and catchment environments. *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology* **251**, 57-82.
- Walker, D., Head, M.J., Hancock, G.J., Murray, A.S., 2000. Establishing a chronology for the last 1000 years of laminated sediment accumulation at Lake Barrine, a tropical upland maar lake, northeastern Australia. *The Holocene* **10**, 415-427.
- Walker, D., Owen, J.A.K., 1999. The characteristics and source of laminated mud at Lake Barrine, northeast Australia. *Quaternary Science Reviews* **18**, 1597-1624.
- Wallbrink, P.J., Hancock, G.J., Olley, J.M., Hughes, A., Prosser, I.P., Hunt, D., Rooney, G., Coleman, R., Stevenson, R., 2003. The Western Port sediment study. CSIRO Land and Water Consultancy Report, Canberra.
- Walsh, R., Shiel, R., Tyler, P., 2001. Reconnaissance Limnology of Tasmania VII. Coastal Lagoons of Bass Strait Islands, with Reference to Endemic Microflora and Microfauna. *Archiv für Hydrobiologie* **152**, 489-510.
- Walsh, R., Shiel, R., Tyler, P., 2004. Reconnaissance Limnology of Tasmania VIII. Tasmanian Coastal Lagoons Epicentres of Endemism in the Australian Aquatic Microbiota. *Proceedings of the Royal Society of Tasmania* **138**, 67-76.
- Walsh, R., Tyler, P., 1998. Reconnaissance Limnology of Tasmania IV. The Distribution and Ecological Preferences of Tasmanian Species of Freshwater Calanoid Copepods (Crustacea:Centropagidae). *Archiv für Hydrobiologie* **141**, 403-420.
- Walsh, R., Tyler, P., 2000. Reconnaissance limnology of Tasmania V. Anthropogenic factors in the distribution of freshwater calanoid copepods (Crustacea: Centropagidae) in Tasmanian lakes. *Archiv für Hydrobiologie* **148**, 625-638.
- Walsh, R.G.J., Vyverman, W., P., T., 1995. Reconnaissance Limnology of Tasmania. III. Coastal lagoons of Bruny Island. *Archiv für Hydrobiologie* **136**, 247-260.
- Wang, X., van der Kaars, S., Kershaw, P., Bird, M., Jansen, F., 1999. A record of fire, vegetation and climate through the last three glacial cycles from Lombok Ridge core G6-4, eastern Indian Ocean, Indonesia. *Palaeogeography, Palaeoclimatology, Palaeoecology* **147**, 241-256.

Wasson, R.J., 1979. Sedimentation history of the Mundi Mundi alluvial fans, western New South Wales. *Sedimentary Geology* **22**, 21-51.

Wasson, R.J., Clark, R.L., Nanninga, P.M., Waters, J., 1987. 210Pb as a chronometer and tracer, Burrinjuck Reservoir, Australia. *Earth Surface Processes and Landforms* **12**, 399 - 414.

Wasson, R.J., Galloway, R.W., 1986. Sediment yield in the Barrier Range before and after European settlement. *The Australian Rangeland Journal* **8**, 79-90.

Watson, J.R., 1995. A Palaeoenvironmental Reconstruction and Assessment of Human Impact in a Supalpine Area of the Snowy Mountains, New South Wales, Australia, University of New South Wales, Sydney.

Webb, 2004. Honours Thesis, University of New South Wales, Sydney.

Whinam, J., Kirkpatrick, J.B., 1994. The Mount Wellington String Bog, Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* **128**, 63-68.

White, C., 6 (9): 426-431., 1967. The prehistory of the Kakadu people. *Mankind* **6**, 426-431.

White, J.P., 1994. Site 820 and the evidence for early occupation in Australia. *Quaternary Australasia* **12**, 21-33.

Williams, J.M., 1980. Crystal Bog, Mount Buffalo: A palaeoecological study of a subalpine peat deposit. Department of Geography, Monash University, Melbourne.

Williams, J.M., 1982. The palaeoecology of a subalpine peat deposit. MA Thesis, Monash University, Melbourne.

Williams, N.J., 2005. The environmental reconstruction of the last glacial cycle at Redhead Lagoon, coastal eastern Australia. PhD Thesis, University of Sydney, Sydney.

Williams, N.J., Harle, K.J., Gale, S.J., Heijnis, H., 2006. The vegetation history of the last glacial—interglacial cycle in eastern New South Wales, Australia. *Journal of Quaternary Science* **21**, 735-750.

Wills, N., 2001. A Palaeolimnological analysis of Lake Cudgen, northern NSW: directions for future management. School of Environmental Science and Management, Southern Cross University, Lismore.

Witt, G.B., Luly, J., Fairfax, R.J., 2006. How the west was once: vegetation change in south-west Queensland from 1930 to 1995. *Journal of Biogeography* **33**, 1585-1596.

Yezdani, G.H., 1970. A study of the Quaternary vegetation history in the volcanic lakes region of western Victoria. PhD Thesis, Monash University, Melbourne.